

Use and usefulness of measures of marine endemism in South Africa

AUTHORS:

Charles L. Griffiths¹

Tamara B. Robinson²

AFFILIATIONS:

¹Department of Biological Sciences and Marine Research Institute, University of Cape Town, Cape Town, South Africa

²Department of Botany and Zoology, Stellenbosch University, Stellenbosch, South Africa

CORRESPONDENCE TO:

Charles Griffiths

EMAIL:

Charles.Griffiths@uct.ac.za

POSTAL ADDRESS:

Department of Biological Sciences and Marine Research Institute, University of Cape Town, Private Bag X3, Rondebosch 7701, Cape Town, South Africa

DATES:

Received: 07 July 2015

Revised: 02 Oct. 2015

Accepted: 08 Oct. 2015

KEYWORDS:

Endemic species; range restriction; biodiversity; marine conservation; taxonomy

HOW TO CITE:

Griffiths CL, Robinson TB. Use and usefulness of measures of marine endemism in South Africa. *S Afr J Sci.* 2016;112(3/4), Art. #2015-0249, 7 pages. <http://dx.doi.org/10.17159/sajs.2016/20150249>

Numerous authors have cited numbers, or proportions, of endemic species within South(ern) African marine taxa, but comparisons between these statistics are confounded by differing definitions of regional boundaries and differences among data sets analysed. These have resulted in considerable variations in published endemism data, even within the same taxonomic group. We tabulated and compared key endemism statistics for regional marine taxa and explained biases in the data sets. The most comprehensive data sets available give overall marine endemism within the national boundaries of South Africa as 28–33%, but estimates within individual taxa making up these totals vary enormously, from 0% (Aves, Mammalia) to over 90% (Polyplacophora). We also examined published data documenting localised endemism patterns around the coastline. These consistently show the highest numbers of endemics occurring along the South Coast. There are logical biogeographical reasons to expect this trend, but endemism rates are also inherently biased by distance from defined political boundaries and by differing sampling effort locally and in neighbouring countries. Range restriction is considered a better measure of conservation status than endemism, although it is far less often used and yields very different patterns. Properly and consistently calculated measures of national endemism do, however, retain significant conservation value, and the rates for South African marine biota are high relative to other regions globally, being exceeded only by New Zealand and Antarctica. It is important that when citing endemism statistics, researchers and conservation managers understand the definitions used and the many constraints under which these measures are derived.

Introduction

Endemism is a widely employed concept, used both as a proxy for range restriction, and as an indicator of the uniqueness and conservation value of the fauna and/or flora within a particular geographical (usually politically defined) region. For example, high proportions of endemism within various South African taxa have been cited in support of the region being recognised as a 'biodiversity hotspot of international importance'.¹ Despite its wide use, endemism is, however, a loosely defined concept – the usual dictionary definition being some variation of 'native or limited to a certain region'.

In practice, the most commonly used boundaries for 'the region' are national political borders, hence the most common regional usage is 'endemic to South Africa'. However, because authors are free to define their own boundaries, these can extend to incorporate wider politically defined regions ('southern Africa'), or a region defined by latitudinal boundaries ('Africa South of 20S') – see marine examples below. Many authors also report rates of endemism for sub-regions within the country, both for biomes, such as the 'thicket biome'², 'succulent karoo' or 'fynbos'³, or for narrower regions, such as 'north-eastern Transvaal Escarpment'⁴, 'southern Langeberg Mountains'⁵, or 'the Cape Peninsula'⁶. In these cases, the numbers reported are of species globally unique to that biome or site (i.e. endemic to the defined region).

Another group of authors divides the region into grid squares^{7,8}, or the coastline into 50 km or 100 km units^{9,10}, then report on the numbers of endemic species present in each square or unit. It is important to note that what is then reported is numbers of species endemic to South Africa that are found in each square/unit (i.e. the same endemic species are reported for multiple squares or units). A few marine authors^{10,11} do subsequently plot numbers of species unique or endemic to only one to three coastal units ('range restricted endemics'). These two methods, of course, yield very different numbers of endemics and geographical patterns of endemism.

The marine fauna of South Africa is no exception with regard to these confusions, and many authors have published endemism data for various taxa in the region (see Table 1), but have used a variety of definitions of endemism, greatly complicating the interpretation of these data. The most commonly used definition, and that most comparable to data from other countries, is that of species restricted to the exclusive economic zone (EEZ) of continental South Africa. However, given the continuity of the South African EEZ with those of adjacent countries, the important role South African researchers have historically played in taxonomic research in the wider region, and other political considerations (e.g. Namibia having long been a protectorate of South Africa), several other definitions have been used. These include latitudinal definitions ('Africa south of 20S' or 'Africa south of the Tropic of Capricorn'), or wider definitions of a 'southern African' region – most often including the whole of Namibia, but only parts of Mozambique. Some studies also restricted their analyses to some maximum depth, while others included the entire EEZ. Clearly, each of these definitions results in different numerical measures of endemism, even when based on the same data sets.

Table 1: Measures of endemism for selected marine taxa from southern Africa. Taxa selected are 'major' grouping for which at least two separate endemism estimates are available

Group	Reference	Region considered	Number of species	Number of endemic species	% endemic species
Mammalia	12	SA EEZ	43	0	0
	13	SA EEZ	44	0	0
Aves	12	SA EEZ	222	0	0
	13	SA EEZ	222	0	0
Reptilia	12	SA EEZ	6	0	0
	13	SA EEZ	6	0	0
Pisces	12	SA EEZ	2000	280	14
	9	SA <200m	1239	101	8
	13	SA EEZ	2000	280	14
Ascidiacea	10	SA <100m	134	72	54
	12	SA EEZ	140	81	58
	14	Namibia-central Mozambique; <15m	135	85	63
	15	Angola-southern Mozambique including islands	168	79	47
	13	SA EEZ	145	45	31
Echinodermata	16	Africa south of 21.5°S	280	134	48
	17	Africa south of 21.5°S	407	190	47
	12	SA EEZ	410	187	46
	10	SA; <100m	65	12	19
	14	Namibia-central Mozambique; <15m	150	65	43
Platyhelminthes	12 (incorrectly summed as 28 with 17 endemic in that paper)	SA EEZ	56	34	61
	13	SA EEZ	254	17	7
Nematoda	12	SA EEZ	338	30	9
	13	SA EEZ	338	30	9
Copepoda	12	SA EEZ	429	41	10
	13	SA EEZ	429	41	10
Cirripedia	12	SA EEZ	86	23	27
	13	SA EEZ	86	23	27
	18	SA EEZ	86	22	25
Amphipoda	19	Africa south of 20°S	292	134	46
	12	SA EEZ	454	149	33
	10	SA <100m	194	77	40
	14	Namibia-central Mozambique; <15m	189	77	41
	13	SA EEZ	454	149	33
Cumacea	20	Africa south of 20°S	75	67	89
	12	SA EEZ	98	70	71

Group	Reference	Region considered	Number of species	Number of endemic species	% endemic species
	13	SA EEZ	98	70	71
Isopoda	12	SA EEZ	300	255	85
	10	SA <100m	252	224	84
	14	Namibia-central Mozambique <15m	278	231	83
	13	SA EEZ	300	255	85
Decapoda	12	SA EEZ	750	150	20
	13	SA EEZ	750	150	20
Polychaeta	12	SA EEZ	760	160	21
	10	SA <100m	523	108	21
	14	Namibia-central Mozambique <15m	556	170	31
	13	SA EEZ	760	161	21
Polyplacophora	12	SA EEZ	29	26	90
	10	SA <100m	23	18	78
	14	Namibia-central Mozambique <15m	24	22	92
	13	SA EEZ	29	26	90
Gastropoda	12	SA EEZ	2262	1258	56
	13	SA EEZ	2262	1258	56
(subgroups combined)	10	SA <100m	998	552	55
	14	Namibia-central Mozambique <15m	919	575	63
Bivalvia	12	SA EEZ	560	270	48
	10	SA <100m	208	93	45
	14	Namibia-central Mozambique <15m	200	89	45
	13	SA EEZ	650 [†]	270	42
Bryozoa	12	SA EEZ	280	99	34
	13	SA EEZ	270	172	63
Actinaria	12	SA EEZ	43	19	44
	21	SA EEZ	47	23	49
	13	SA EEZ	43	19	44
	22	SA EEZ	61	28	46
Octocorallia	12	SA EEZ	204	110	54
	10	SA <100m	54	16	30
	14	Namibia-central Mozambique <15m	68	55	81
	13	SA EEZ	204	110	54
Hydrozoa	12	SA EEZ	457	86	19
	13	SA EEZ	382	86	23
Porifera	12	SA EEZ	289	10	3
	13	SA EEZ	346	32	9
Combined fauna	12	SA EEZ	11 130	3496	31
	10	SA; <100m	2533	931	26

Group	Reference	Region considered	Number of species	Number of endemic species	% endemic species
	14	Namibia-central Mozambique <15m	2650	1401	53
	13	SA EEZ	12 914	4233	33
	23	SA EEZ	12715	3549	28

[†]Probably transposed incorrectly from 560 of Gibbons et al.¹²

For more complete taxonomic listings, see Gibbons et al.¹² (Table 1) and Griffiths et al.¹³ (Table S1). Geographic region and depth zones are specified for each estimate (EEZ = Exclusive Economic Zone).

Actual levels of endemism are also subject to constant change over time in response to developments in taxonomic knowledge, both within and outside of the region under consideration. Simply put, new taxonomic data from within the study region will *increase* endemism when new species are described from within the study area, but will *reduce* endemism when species already known elsewhere are newly reported there. Conversely, studies done *outside* of the study area will depress endemism levels if they record species previously thought to be endemic to South(ern) Africa. Keeping track of these changes requires a high level of taxonomic expertise and many regional publications have in fact based their analysis on historical data, derived from dated taxonomic monographs or museum catalogues, without correcting these to take account of more recent taxonomic revisions and discoveries, particularly those carried out *outside* of the study region. Such estimates must thus be treated with caution.

Another group of studies have analysed how the numbers, or proportions, of endemic marine species vary around a series of equal 50 km or 100 km units distributed around the length of the South(ern) African coastline. Such studies include those on fish⁴ and on various invertebrate taxa.^{10,11,13} In addition to the biases and constraints already mentioned, there is also an inherent bias in such 'within region' analyses, in that such levels of endemism are partially determined by distance away from the defined boundary. In other words, if all species had randomly-centred 500 km ranges, all species from sites more than 500 km from the defined boundary would by necessity be considered endemic. In this context, measures of range-restriction may be more indicative of rarity or conservation status than the far more commonly-used endemism, although they too suffer from biases related to sampling effort, as discussed below.

However flawed they are in biological terms, endemism statistics do still have conservation value, as they do give an indication of the number (or proportion) of described species that fall under the responsibility of a particular national government or conservation agency, or of the proportions that are conserved within existing protected areas¹⁰. Thus a critical appraisal of the use of endemism statistics is necessary if we are to ensure that prioritisation of conservation goals is underpinned by sound scientific inputs, of which endemism is an important example.

The South African marine environment provides a particularly good, and unusual, model system to undertake such an analysis. Firstly, a large number of regional endemism measures have been published and (unusually in the global context) a variety of definitions have been applied to delimit the region under consideration. The South(ern) African marine environment is also a good model to analyse in that the coastline is almost linear, with very few inlets, bays or islands, plus almost all samples have been collected within a fairly narrow band adjacent to that coastline.¹³ This means that it is easy to re-analyse the data in terms of linear distance from the boundary, or in terms of linear measures of range-restriction.

In this paper we aim to explore the use and usefulness of the concept of endemism, as applied to the marine fauna of South(ern) Africa. We do this in several ways: firstly, by reviewing previous studies that have calculated endemism and comparing the data so obtained in the light of

the definitions and data sources used and, secondly by discussing factors that affect reported levels of endemism. Lastly, we compare measured marine endemism statistics in South Africa with those from other comparable regions and debate the usefulness of alternative measures.


Existing measures of endemism

Endemism measures have been published for a large variety of South(ern) African marine taxa, of which we have reproduced only a selected subset of 'major' groups, especially focussing on those for which multiple estimates are available (Table 1). The tremendous variation in absolute measures of endemism (from zero in several vertebrate taxa, to over 80% in Cumacea, Isopoda and Polyplacophora) is notable. There is also considerable variability between estimates made for the same taxonomic group, but by different authors. These may have resulted from inclusion of different data sets, but most obviously vary with the definition of the study area, with larger study areas generally resulting in the inclusion of the ranges of more species and thus in higher endemism levels. For some taxa the various estimates are identical, as either the exact same data set has been analysed, or because later authors have cited figures directly from earlier ones, without any new analyses. In both cases, inaccuracies are likely to result, as in reality, new research both inside and outside the study area will have resulted in changing proportions of endemics, as discussed above.

Comparing groups, one would expect pelagic taxa to have much lower endemism levels than benthic ones, as pelagic taxa are far more mobile. This certainly applies to some taxa, for example, the Copepoda, the majority of which are pelagic, have much lower endemism figures than most benthic invertebrate groups. Indeed, if pelagic taxa within the Copepoda are separated from benthic ones, this trend becomes dramatic – pelagic copepods being only 1% endemic and benthic ones 49% endemic.¹² The same is true within the Amphipoda, which also contain both pelagic taxa (Hyperideae), which are only 1% endemic, and benthic taxa (Gammarideae and Caprellideae), which together show 45% endemism.¹² One might also expect that broadcast spawning taxa might have lower endemism levels than brooding ones, but clear distinctions among benthic taxa, based on life history, are not apparent within these data sets. For example, while endemism levels among some brooding taxa, such as Cumacea (71–89%) and Isopoda (83–85%) are, as expected, exceptionally high, equally high figures are evident among (mainly) broadcast spawning groups, such as Polyplacophora (78–92%) (Table 1). Conversely, Echinodermata (generally broadcast spawners) have lower endemism ratios (19–48%), similar to the brooding Amphipoda (33–46%).

The states of taxonomic knowledge, both within the region of study and in adjoining areas, have major impacts on reported levels of endemism. Clearly, if few species have been described from within a study region, there can be few endemic species reported from that region. Thus one would expect taxa that have been poorly studied within South(ern) Africa to have artificially depressed levels of reported endemism. This is true, for example, for Nematoda at 9%. Conversely, for well-studied taxa, most of the actual endemic species present would have been discovered and described and hence endemism ratios should be higher (and more realistic). These expectations are validated when plotting the relationship between numbers of endemic species and the state

of knowledge of various marine taxa for South Africa (Figure 1). Note that this relationship breaks down for the few taxa in the highest state of knowledge category, simply because these include only vertebrate groups and mangroves, which comprise large, highly mobile species with large ranges, and brown algae, for which no endemism data are available.¹³ While large size seems to equate with low endemism in those taxa, as is also reported by Costello et al.²⁴, only a very weak positive correlation between size and range was observed among South African invertebrate taxa.¹⁴



Source: Table S1 in Griffiths et al.¹³

Figure 1: The proportions of endemic species in various taxonomic groups relative to their states of taxonomic knowledge as rated from poor (1) to excellent (5).

Additional biases in reported levels of endemism are also created by state of knowledge in adjoining regions. Thus if the fauna, or a particular taxonomic group, is well known within the study area, but poorly known in adjoining regions, this artificially inflates the reported endemism ratio, because species whose real ranges extend into adjoining regions are unlikely to have been detected there. Conversely, if the fauna in adjoining regions is well known, then species with broad ranges will most likely have been detected there, and endemism within the study area will be lower (and more realistic).

An interesting example of this is the comparison between the biologically similar and closely related Amphipoda and Isopoda (Table 1). The former are well studied in Madagascar^{25,26} and have a relatively low endemism ratio, while the Isopoda, which are well known in South Africa,²⁷ but poorly studied elsewhere in Africa, have a much higher apparent proportion of 'endemic' species.

The effects of increased study area on the overall endemism ratio of the fauna as a whole is particularly clear, with Gibbons et al.¹², Awad et al.¹⁰, Costello et al.²³ and Griffiths et al.¹³ giving similar overall marine endemism estimates for South Africa (31%, 26%, 28% and 33% respectively), while Scott¹⁴ gives a much higher estimate of 53% for the inclusive southern African region (Namibia to central Mozambique). Note, however, that even this comparison is not based on identical data sets, as each of these studies included a somewhat different number and mix of taxa, and both Awad et al.¹⁰ and Scott¹⁴ considered only coastal fauna, whereas Gibbons et al.¹², Costello et al.²³ and Griffiths et al.¹³ examined the entire EEZ (and derived much of their data from the same sources). When we look at individual taxa, the trend remains for higher endemism whenever a wider area is examined. For example, 55–56% endemism for South African Gastropoda vs 63% for the wider southern African region; 21% of South African Polychaeta vs 31–36% for the wider region; 71% vs 89% for Cumacea, etc. There are, however, exceptions, such as Primo & Vazquez¹⁵, who give a lower endemism estimate of 47% for the Ascidiacea of the wider Angola–Mozambique region than either Gibbons et al.¹² or Awad et al.¹⁰ (but not Griffiths et al.)¹³ provide for South Africa alone (58%, 54% and 31% respectively). This can only be the result of different data sets being used in these analyses.

Several studies have reported on geographical patterns of endemism within the region,^{9,10,11} plotting patterns of both species richness and endemism within particular taxa as one moves around the coastline of South(ern) Africa (Figure 2). The usual findings are that numbers (and proportions) of endemic species peak along the South Coast. There are good biological reasons to expect this trend, in that the warm temperate South Coast is both situated exclusively within the political boundaries of South Africa and is geographically isolated from other warm temperate regions (both features which promote high endemism). By contrast, the cool temperate west coast and more tropical east coast bioregions are contiguous with similar climatic regions in adjacent countries, meaning that species living in those habitats are likely to range across political borders and hence not be endemic.




Figure 2: (a) Numbers of range-restricted (range < 300 km) species/100 km stretch of coastline and (b) marine invertebrate species richness/100 km. Distance around coast measured from Namibian border to Mozambique border. Data from Awad et al.¹⁰ Ecoregions described by Sink et al.²⁸ are indicated in panel (a) as SB (Southern Benguela), AG (Agulhas), N (Natal) and DG (Delagoa). The dark sections of the bars in panel (b) indicate endemic species.

It is important to note here, though, that endemism is strongly influenced simply by distance from the political borders of South Africa and is not a good indicator of either range restriction or rarity. For example, a species with a short range that happens to straddle a political border will not be endemic to either country, but may have a very restricted range and hence require conservation attention. Conversely, one that is endemic can have a much larger range (of up to 3000 km in this case) and hence be of less conservation concern. Endemism along the South Coast of South Africa is thus inflated simply because this region is furthest from the political borders used to define endemism in this region.

Marine endemism rates in South Africa in the global context

Table 2 shows known rates of marine endemism in a variety of global regions, as summarised by Costello et al.²³ It is notable that South Africa supports the third highest proportion of endemic species among listed regions, despite having one of the smaller seabed areas and supporting comparatively few described species, relative to some other listed regions. The reasons for these patterns are complex and involve not only 'real' factors, such as the size, isolation and taxonomic uniqueness of the fauna of regions under consideration, but artefacts associated with sampling and taxonomic effort, both in the region in question and in neighbouring areas.

Table 2: Numbers of marine species reported from various regions globally, the percentage of these that are endemic and size of region.

Region	Total number of species	% endemics	Seabed area km ²
Antarctica	8200	45	21 186 153
Australia	32889	28	6 819 501
Baltic	5865	2	411 218
Caribbean	12046	13	2 828 125
China	22365	7	831 966
Japan	32777	6	3 970 743
Mediterranean	16845	7	2 451 059
New Zealand	12780	51	4 073 895
South Africa	12715	28	846 463

Source: Costello et al.²³

Regions that are large and/or isolated (Australia, New Zealand, Antarctica) would be expected to, and do, have high levels of endemism. Conversely, those with relatively small areas and/or which lie adjacent to other well-studied areas (China, Japan, Baltic, Mediterranean, Caribbean) have relatively low endemism. South Africa might appear to be an outlier here, as it has a relatively small sea area, but a high endemism ratio. However, it does incorporate an 'island' of warm temperate water that is very isolated geographically, from other warm temperate regions, and which hosts the majority of the endemic species. It is also something of an 'island of taxonomic expertise' in a much wider African region, within which marine biodiversity is poorly documented, and this tends to (artificially) elevate rates of endemism.

Is range restriction a better measure than endemism?

The problems outlined above, with respect to accurate determination of endemism rates, beg the question whether endemism is a useful concept for managers. Clearly, it does have political importance in that the conservation of endemic species falls exclusively under the jurisdiction of a particular nation's conservation authorities. It is also a term the public associate with and which has implications for funding of conservation efforts through eco-tourism, etc.

On the other hand, range restriction and population size or rarity (which are not dealt with here, are more difficult to measure and more rarely reported in marine taxa) are much more important concepts in terms of viability and vulnerability of the population of a given species. Studies by Awad et al.¹⁰ and Scott et al.¹¹ have plotted the geographic distributions of range-restricted species along the South Africa coast, showing range-restricted forms to be strongly localised relative to endemic ones. Specifically, range-

restricted marine species tend to be strongly concentrated at focal sites in the vicinities of Cape Point, Port Elizabeth and to a lesser extent, Durban (Figure 2a). Scott et al.¹¹ suggest that, as these sites tend to correspond with biogeographic breaks, these peaks may represent species living at the ecotone between biogeographic provinces. However, these specific locations coincidentally also fall at the locations of the three major coastal centres of marine research in the country and are, therefore, also sites of enhanced sampling effort and species discovery. As these well-sampled areas thus contain 'extra' species not reported in adjacent sectors of coast (Figure 2), it is thus inevitable that these 'extra' species will appear to be range-restricted to those sites. Moreover, while the intensely-sampled area around Cape Point does indeed coincide exactly with the boundary between the Southern Benguela and Agulhas ecoregions,²⁸ the peaks around Port Elizabeth and Durban are offset from biogeographic breaks by at least 300 km. This strongly suggests that at least these peaks in locally endemic species are largely a function of differences in sampling effort, rather than true range restriction. The larger peak around Cape Point may be more 'real' as it is situated in a region with unique ecotonal characteristics, but the greatly enhanced sampling effort at this site is certainly responsible for at least some of these apparently 'localised' species.

Acknowledgements

Financial support for this research was provided by a National Research Foundation grant to CLG, through the SeaKeys programme.

Authors' Contributions

The paper was a joint effort by C.L.G. and T.B.R. C.L.G. conceived the idea for the paper and wrote the original draft. T.B.R. edited and added to the draft and drew the figures. Being a review, there were no original experiments, only data extraction and interpretation which we did together.

References

1. Endangered Wildlife Trust. The Biodiversity of South Africa 2002: Indicators, trends and human impacts. Cape Town: Struik Publishers; 2002.
2. Vlok JHJ, Euston-Brown DIW, Cowling RM. Acocks' Valley bushveld 50 years on: New perspectives on the delimitation, characterisation and origin of subtropical thicket vegetation. *S Afr J Bot.* 2003;69:27–51. [http://dx.doi.org/10.1016/S0254-6299\(15\)30358-6](http://dx.doi.org/10.1016/S0254-6299(15)30358-6)
3. Verboom GA, Archibald JK, Bakker FT, Bellstedt DU, Conrad F, Dreyer LL, et al. Origin and diversification of the Greater Cape flora: Ancient species repository, hot-bed of recent radiation, or both? *Molec Phylogenetics Evol.* 2009;51:144–53. <http://dx.doi.org/10.1016/j.ympev.2008.01.037>
4. Matthews WS, Van Wyk AE, Bredenkamp GJ. Endemic flora of the north-eastern Transvaal escarpment, South Africa. *Biol Conserv.* 1993;63:83–94. [http://dx.doi.org/10.1016/0006-3207\(93\)90077-E](http://dx.doi.org/10.1016/0006-3207(93)90077-E)
5. McDonald DJ, Cowling RM. Towards a profile of an endemic mountain fynbos flora: Implications for conservation. *Biol Conserv.* 1995;72:1–12. [http://dx.doi.org/10.1016/0006-3207\(94\)00030-T](http://dx.doi.org/10.1016/0006-3207(94)00030-T)
6. Picker, MD, Samways MJ. Faunal diversity and endemism of the Cape Peninsula, South Africa – a first assessment. *Biodiver Conserv.* 1996;5:591–606. <http://dx.doi.org/10.1007/BF00137611>
7. Skelton PH, Cambray JA, Lombard AT, Benn GA. Patterns of distribution and conservation of freshwater fishes in South Africa. *S Afr J Zool.* 1995;30:71–81. <http://dx.doi.org/10.1080/02541858.1995.11448375>
8. Drinkrow DR, Cherry MI. Anuran distribution, diversity and conservation in South Africa, Lesotho and Swaziland. *S Afr J Zool.* 1995;30:82–90. <http://dx.doi.org/10.1080/02541858.1995.11448376>
9. Turpie JK, Beckley LE, Katua SM. Biogeography and the selection of priority areas for conservation of South African coastal fish. *Biol Conserv.* 2000;92:59–72. [http://dx.doi.org/10.1016/S0006-3207\(99\)00063-4](http://dx.doi.org/10.1016/S0006-3207(99)00063-4)
10. Awad AA, Griffiths CL, Turpie JK. Distribution of South African marine invertebrates applied to the selection of priority conservation areas. *Divers Distrib.* 2002;8:129–145. <http://dx.doi.org/10.1046/j.1472-4642.2002.00132.x>
11. Scott RJ, Griffiths CL, Robinson TB. Patterns of endemism and range-restriction among southern African coastal marine invertebrates. *Afr J Mar Sci.* 2012;34:341–348. <http://dx.doi.org/10.2989/1814232X.2012.725284>
12. Gibbons MJ. The taxonomic richness of South Africa's marine fauna - a crisis at hand. *S Afr J Sci.* 1999;95:8–12.

13. Griffiths CL, Robinson TB, Lange L, Mead A. Marine biodiversity in South Africa – state of knowledge, spatial patterns and threats. *PLoS ONE* 2010;5(8):e123008. <http://dx.doi.org/10.1371/journal.pone.0012008>
14. Scott R. 2009. Biogeographical patterns in southern African marine invertebrates [MSc dissertation]. Cape Town: University of Cape Town; 2009.
15. Primo C, Vázquez E. Zoogeography of the southern African ascidian fauna. *J Biogeogr.* 2004;31:1987–2009. <http://dx.doi.org/10.1111/j.1365-2699.2004.01144.x>
16. Clark AM, Courtman-Stock J. *The Echinoderms of Southern Africa*. London: British Museum (Natural History); 1976.
17. Thander AS. Zoogeography of the southern African echinoderm fauna. *S Afr J Zool.* 1989;24:311–318.
18. Biccard A. Taxonomy, systematics and biogeography of South African Cirrepedia (Thoracica) [MSc dissertation]. Cape Town: University of Cape Town; 2012.
19. Griffiths CL. *The gammaridean and caprellid Amphipoda of southern Africa* [PhD thesis]. Cape Town: University of Cape Town; 1974.
20. Day JA. *Southern African Cumacea* [PhD thesis]. Cape Town: University of Cape Town; 1978.
21. Acuña FH, Griffiths CL. Species richness, endemism and distribution patterns of South African sea anemones (Cnidaria: Actiniaria & Corallimorpharia). *Afr Zool.* 2004;39(2):193–200.
22. Laird CM. Taxonomy, systematics and biogeography of South African Actiniaria and Corallimorpharia [PhD thesis]. Cape Town: University of Cape Town; 2013.
23. Costello MJ, Coll M, Danovaro R, Halpin P, Ojaveer H, Miloslavich P. A census of marine biodiversity knowledge, resources and future challenges. *PLoS ONE* 2010;5(8):e12110. <http://dx.doi.org/10.1371/journal.pone.0012110>
24. Costello MJ, May RM, Stork NE. Can we name Earth's species before they go extinct? *Science.* 2013;339(6118):413–416. <http://dx.doi.org/10.1126/science.1230318>
25. Ledoyer M. Crustacés amphipodes gammariens. Familles des Acanthonotozomatidae à Gammaridae [Gammaridean amphipod crustaceans. Familles Acanthonotozomatidae to Gammaridae]. *Faune de Madagascar.* 1982;59:1–598. French.
26. Ledoyer M. Crustacés amphipodes gammariens. Familles des Haustoriidae à Vitjazianidae [Gammaridean amphipod crustaceans. Familles Haustoriidae to Vitjazianidae]. *Faune de Madagascar.* 1986;59:599–1112. French.
27. Kensley B. *Guide to the marine isopods of Southern Africa*. Cape Town: Trustees of the South African Museum; 1978.
28. Sink K, Holness S, Harris L, Majiedt P, Atkinson L, Robinson T, et al. *National Biodiversity Assessment 2011: Technical Report. Volume 4: Marine and Coastal Component*. Pretoria: South African National Biodiversity Institute; 2012.

